
Coronal signatures of flares and CMEs

Dr. Karin Muglach NASA/GSFC and Artep, Inc.

SW-REDI 2015



Chromosphere





Large Scale Structures Near the Solar Surface

two kinds of measurement to collect information about the Sun:

Remote Sensing and In-situ Measurement

Key for remote sensing of the sun (and stars): Solar Spectrum








complete solar spectrum and EUV part of solar spectrum

Key for remote sensing of the sun (and stars): Solar Spectrum


True-Color Irradiance Spectrum 392 to 692 nm from Kitt Peak Residual Irradiance Atlas (Kurucz 2005)







Global magnetic field (extrapolation): 3d structure Line-of-sight full disk magnetogram: 2d cut at photosphere




Full disk white light image (SDO), full disk line-of-sight magnetogram (SDO)

Active Region evolution in white light and magnetogram (SDO).


If we just have white light images and magnetograms: Q: How are the polarities connected?

If we just have white light images and magnetograms: Q: How are the polarities connected?


- A1: extrapolation
- A2: corona images: outline (some) of the magnetic field connectivity!



Full disk magnetogram and 171 image (SDO)




Full disk image in H alpha (BBSO): filaments seen as dark absorption structures




High resolution image in H alpha (Dutch Open Telescope) filaments seen as dark absorption structures

Example of filaments:

- Quiescent filament in high spatial resolution (Hinode SOT)
- Filament eruption (SDO, composite)



SIMPLE (!!) cartoon of filament magnetic field



magnetic field lines almost parallel to PIL

Notes on filaments:


- Filament: on-disk structure (seen in absorption)
 Prominence: same structure off limb (seen in emission)
- Best wavelengths: H alpha, He II 304, Fe XII 195 A (AIA, STEREO)
- All filaments have a PIL
- But not all PILs are filaments!
- Caution: full disk magnetograms give only the line-of-sight magnetic field – projection effects near the solar limb!

Solar Eruptions: Flares and CMEs

- Energy is stored in the solar magnetic field (active regions and filaments): accumulated over a long period of time – days, weeks, months
- Energy is released in eruptions (flares, CMEs): in a short time scale (minutes, hours)

Magnetic energy is converted to thermal energy (and radiative energy) and kinetic energy (e.g. mass motion in CMEs and SEPs)

Solar Eruptions: Flares and CMEs



one possible scenario for an eruption:

- reconnection at the x-point (energy release)
- CME escapes upward, field-lines open up
- Post-eruptive loops appear below x-point (additional heating)

Solar Eruptions: Flares and CMEs

Caution: the real sun is more complicated compared to the cartoon – e.g. magnetic field is a

3d structure

- some eruptions show no/very little X-ray signature (particularly filament eruptions)
- some flares have no CMEs

Large scale structures in the corona

- Images: SDO AIA 193 A, STEREO EUVI 195 A (filter contains Fe XII 195 A line, T~1.5 MK)
- Line-of-sight magnetograms: polarity inversion line (PIL)
- Active Regions: bi-polar, bright (emission), closed magnetic field (field lines perpendicular to PIL)
- Filaments: bi-polar, dark (absorption), closed magnetic field (field lines parallel to PIL)
- Coronal hole: uni-polar, dark (less dense), open magnetic field

Coronal signatures of CMEs

- Data to use: SDO AIA, STEREO EUVI (A & B)
- Brightenings: flares, post-eruptive arcade (193), arcade footpoints (304, 193)
- Darkenings: dimmings (transient coronal holes), dark/absorbing/cool material rising (filament eruption)
- Off-limb: opening of closed coronal field lines, AIA 304 emission structure
- Not a signature of eruption: active region loop brightenings, (small) flares

Coronal signatures of CMEs

Good period to study: SDO 2014-02-18 - 21 (use AIA 211, 193, 304)