


Space Weather from

Coronal Holes and High Speed Streams



Karin Muglach (NASA/GSFC and CUA)

Based on a presentation by
M. Leila Mays (NASA/GSFC and CUA)

SW REDI

June 2016




The **solar wind** is a flow of **plasma** and the frozen-in solar **magnetic field** from the Sun. The outward flow is due to the gas pressure difference between interplanetary space and the solar corona.

Changes in the solar magnetic field (from solar activity) influence the solar wind which, in turn, influences planets, spacecraft, and other bodies inside the solar wind (the **heliosphere**).

TABLE 4.1. Observed Properties of the Solar Wind near the Orbit of the Earth (1 AU)

Proton density	6.6 cm^{-3}
Electron density	7.1 cm^{-3}
He ²⁺ density	0.25 cm^{-3}
Flow speed (nearly radial)	$450 \text{ km} \cdot \text{s}^{-1}$
Proton temperature	$1.2 \times 10^5 \text{ K}$
Electron temperature	$1.4 \times 10^5 \text{ K}$
Magnetic field (induction)	$7 \times 10^{-9} \text{ tesla (T)}$

near the Earth, compared to the magnetosphere, the solar wind plasma (mostly ionized Hydrogen) is hot, tenuous, and fast moving, and the weak magnetic field is nearly parallel to the ecliptic plane, but 45° to the Sun-Earth line



Due to solar **rotation** parcels of solar wind plasma leaving the sun form a spiral (analogous to the water spirals formed from a rotating sprinkler), which is called the **Parker spiral**. The angle that a solar wind magnetic field line makes at 1 AU is close to 45 degrees.





FIG. 4.5. Loci of a succession of fluid parcels (eight of them in this sketch) emitted at constant speed from a source fixed on the rotating sun.

$$U_{\phi} = -\omega r$$

Solar wind can be divided into fast and slow wind components.



Solar wind can be divided into fast and slow wind components.

Fast wind

450–800 km/s

$n_p \sim 3 \text{ cm}^{-3}$

~95% H, 5% He, minor ions and same number of electrons

$T_p \sim 2 \times 10^5 \text{ K}$

$B \sim 5 \text{ nT}$

Alfvénic fluctuations

Origin in coronal holes

Slow wind

$< \sim 450 \text{ km/s}$

$n_p \sim 7\text{--}10 \text{ cm}^{-3}$

~94% H, ~4% He, minor ions and same number of electrons – great variability

$T_p \sim 4 \times 10^4 \text{ K}$


$B \sim 4 \text{ nT}$

Density fluctuations


Origin 'above' coronal streamers and through small-scale transients

(Bothmer and Zhukov, 2007)

Coronal holes appear as dark areas on the solar surface in the **EUUV** (extreme ultraviolet) and **X-ray** radiation. They have a **lower density** and **temperature** compared to the surrounding corona. **Coronal holes** correspond to regions of open magnetic fields. Visible best in lines with temperatures more than 1.5 MK.




SDO/AIA- 211 20110425_031014



SDO/AIA 193 2011-04-25 03:28:20 UT

Large **polar coronal holes** are persistent for about 7 years around solar minimum. During solar maximum and high solar activity **coronal holes** exist at all latitudes, but are less persistent.



Coronal holes correspond to regions of open magnetic fields.






FIG. 4.8. Photograph of the corona obtained at the 1966 solar eclipse and a sketch of the magnetic-field structures believed to exist within the observed coronal features.



High speed solar wind streams are formed by higher speed solar wind originating from **coronal holes**. Higher speed streams are less tightly wound in the Parker spiral compared to slower ones, and at various distances the faster solar wind **overtakes** the slower wind ahead of it.




FIG. 4.13. Geometry of the interaction between fast solar wind (on less tightly wound spiral streamlines) and ambient solar wind (on more tightly wound spiral streamlines). The plasma is compressed where streamlines converge. (From Pizzo, 1985.)

A **stream interaction region** (SIR) forms at the compressed boundary between the fast and slow solar wind in a high speed stream. High speed streams from persistent coronal holes over multiple solar rotations are called **corotating interaction regions** (CIRs).




FIG. 4.13. Geometry of the interaction between fast solar wind (on less tightly wound spiral streamlines) and ambient solar wind (on more tightly wound spiral streamlines). The plasma is compressed where streamlines converge. (From Pizzo, 1985.)

Example of an **high speed solar wind stream** observed in-situ at ACE

iSWA layout: <http://go.nasa.gov/17nkicp>

The increase in speed from a solar wind high speed stream pumps energy into the magnetosphere which can cause **geomagnetic storms** and **energizes particles**.


They can produce energetic **electron flux enhancements** in the radiation belt.

Geomagnetic storms are disturbances/changes in Earth's magnetic field due to changes in solar wind conditions typically lasting 3-6 days.

High speed streams can also cause geomagnetic storms, however they are longer in duration and not as strong as geomagnetic storms caused by CMEs.

(the magnetosphere lesson will go into more detail)

A high speed stream can cause an energetic **electron flux enhancement** in the radiation belt.



iSWA layout: <http://go.nasa.gov/17nkicp>

Slide link summary

SW REDI website

<http://ccmc.gsfc.nasa.gov/support/SWREDI/swredi.php>

iSWA <http://iswa.gsfc.nasa.gov>

iSWA Cygnet Glossary http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Full_iSWA_Cygnet_List

iSWA Space Weather Glossary <http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Glossary>

Most figures and tables in the slides are from the “Introduction to Space Physics” textbook

<http://www.cambridge.org/us/knowledge/isbn/item1145043>

iSWA layout of a high speed stream observed at ACE <http://go.nasa.gov/17nkicp>

The slide cover is divided into two main sections. On the left, there is a white vertical strip containing three logos: the NASA logo at the top, a stylized gear logo in the middle, and a logo for the 'COMMUNITY' program at the bottom. The right section is a dark background with a composite image of the sun's corona. The text on the slide is as follows:

Space Weather from
**Coronal Holes and
High Speed Streams**

Karin Muglach (NASA/GSFC and CUA)

Based on a presentation by
M. Leila Mays (NASA/GSFC and CUA)

SW REDI June 2016

This lecture will describe features on the sun called coronal holes, and the high speed streams that result from them. The coronal hole can be seen in this SDO AIA composite image (wavelengths 211, 193, and 171 Angstroms), as the dark purple area. The bright areas are active regions.


The **solar wind** is a flow of **plasma** and the frozen-in solar **magnetic field** from the Sun. The outward flow is due to the gas pressure difference between interplanetary space and the solar corona.

Changes in the solar magnetic field (from solar activity) influence the solar wind which, in turn, influences planets, spacecraft, and other bodies inside the solar wind (the **heliosphere**).

TABLE 4.1. Observed Properties of the Solar Wind near the Orbit of the Earth (1 AU)

Proton density	6.6 cm^{-3}
Electron density	7.1 cm^{-3}
He ²⁺ density	0.25 cm^{-3}
Flow speed (nearly radial)	$450 \text{ km} \cdot \text{s}^{-1}$
Proton temperature	$1.2 \times 10^5 \text{ K}$
Electron temperature	$1.4 \times 10^5 \text{ K}$
Magnetic field (induction)	$7 \times 10^{-9} \text{ tesla (T)}$

near the Earth, compared to the magnetosphere, the solar wind plasma (mostly ionized Hydrogen) is hot, tenuous, and fast moving, and the weak magnetic field is nearly parallel to the ecliptic plane, but 45° to the Sun-Earth line



The **solar wind** is a flow of **plasma** and the frozen-in solar **magnetic field** from the Sun. The outward flow is due to the gas pressure difference between interplanetary space and the solar corona.

Changes in the solar magnetic field (from solar activity) influence the solar wind which, in turn, influences planets, spacecraft, and other bodies inside the solar wind (the **heliosphere**).

Table 4.1 (from Kivelson and Russell, Introduction to Space Physics) shows properties of the solar wind near the Earth. Near the Earth, compared to the magnetosphere, the solar wind plasma (mostly ionized Hydrogen) is hot, tenuous, and fast moving, and the weak magnetic field is nearly parallel to the ecliptic plane, but 45° to the Sun-Earth line.

The graph on the right is OMNI combined solar wind data measured at L1, just ahead of Earth (1 AU), showing a typical timeseries of the solar wind average magnetic field vector (top), velocity in the x direction (Vx, GSE coordinates), proton density, and proton temperature for a period in July 2008.

Due to solar **rotation** parcels of solar wind plasma leaving the sun form a spiral (analogous to the water spirals formed from a rotating sprinkler), which is called the **Parker spiral**. The angle that a solar wind magnetic field line makes at 1 AU is close to 45 degrees.





FIG. 4.5. Loci of a succession of fluid parcels (eight of them in this sketch) emitted at constant speed from a source fixed on the rotating sun.

The **Parker spiral**, is the spiral of Archimedes magnetic geometry of the solar wind due to solar **rotation**. Parcels of solar wind leaving the sun are analogous the water spirals formed from a rotating sprinkler. The angle a solar wind magnetic field line makes at 1 AU is close to 45 degrees.

In Figure 4.5 (from Introduction to Space Physics) you can imagine the solar wind as parcels of plasma being released from the sun in succession (starting with parcel #1). As the sun rotates, the parcels are released from the rotated location, forming a spiral shape (parcels #1 through #8).

Solar wind can be divided into fast and slow wind components.



Solar wind can be divided into fast and slow wind components.

In the top panel you can see Ulysses solar wind speed data plotted radially on top of a typical EUV and coronagraph image of the sun from solar minimum to maximum, then to the most recent solar minimum. In the bottom panel the sunspot number is plotted in black, which is a measure of the solar cycle, and in red the heliospheric current sheet tilt is plotted.

From this graph you can see the solar wind slow and fast components readily distinguishable during solar minimum, with faster solar wind (~700 km/s) from the poles, and slower solar wind (~400 km/s) from the equator. During solar maximum slow and fast solar wind is measured at a variety of latitudes.

Notice that in during solar minimum the current sheet is mostly equatorial but it is a highly inclined near maximum.


Solar wind can be divided into fast and slow wind components.

Fast wind	Slow wind
450–800 km/s	<~450 km/s
$n_p \sim 3 \text{ cm}^{-3}$	$n_p \sim 7\text{--}10 \text{ cm}^{-3}$
~95% H, 5% He, minor ions and same number of electrons	~94% H, ~4% He, minor ions and same number of electrons – great variability
$T_p \sim 2 \times 10^5 \text{ K}$	$T_p \sim 4 \times 10^4 \text{ K}$
$B \sim 5 \text{ nT}$	$B \sim 4 \text{ nT}$
Alfvénic fluctuations	Density fluctuations
Origin in coronal holes	Origin 'above' coronal streamers and through small-scale transients

(Bothmer and Zhukov, 2007)

This table from Bothmer and Zhukov (2007), shows typical solar wind values for the fast and slow components. The fast solar wind has a higher speed, lower density, higher temperature and originates from coronal holes. The slow solar wind has lower speeds, higher densities, lower temperatures and originates above coronal streamers or from small transients.

Coronal holes appear as dark areas on the solar surface in the **EUV** (extreme ultraviolet) and **X-ray** radiation. They have a **lower density** and **temperature** compared to the surrounding corona. **Coronal holes** correspond to regions of open magnetic fields. Visible best in lines with temperatures more than 1.5 MK.




The faster solar wind originates from coronal holes. What are coronal holes? **Coronal**

holes appear as dark areas on the solar surface in the **EUV** (extreme ultraviolet) and **X-ray** radiation. They have a **lower density** and **temperature** compared with the surrounding corona. **Coronal holes** correspond to regions of open magnetic fields. Visible best in lines with temperatures more than 105 K.

These images show the Sun at various wavelengths, coronal holes appear as dark areas in the SDO AIA 211 Å (top left) and 193 Å (top right) images.

In the left image, a PFSS model of the coronal magnetic field is overlaid on an SDO AIA 211 Å image. This model illustrates the closed magnetic field line structure over active regions, and the (locally) open coronal magnetic field lines.

Large **polar coronal holes** are persistent for about 7 years around solar minimum. During solar maximum and high solar activity **coronal holes** exist at all latitudes, but are less persistent.



Large **polar coronal holes** are persistent, several years around solar minimum. During solar maximum and high solar activity they exist at all latitudes, but are less persistent.


The polar coronal holes are visible at the poles of the left image from SOHO EIT 195 Angstroms and the right GOES-14 SXI X-ray image for the same period.

Lower latitude coronal holes are also visible in the center of these images.

2010/06/05 00:48

Coronal holes correspond to regions of open magnetic fields.

FIG. 4.8. Photograph of the corona obtained at the 1966 solar eclipse and a sketch of the magnetic-field structures believed to exist within the observed coronal features.



Coronal holes correspond to regions of (locally) open magnetic fields.

The image on the right shows the PFSS magnetic field model overlay on the SDO AIA 211, 193, 171, composite image. Again this model illustrates the closed magnetic field line structure over active regions, and the (locally) open coronal magnetic field lines.

The top right corona photograph from an eclipse (from Introduction to Space Physics) show coronal helmet streamers as bright structures, and the dark areas correspond to coronal holes. The bottom right schematic for this photograph shows the presumed magnetic field configuration of the corona for this photograph. Helmet streamers appear above closed magnetic field lines and prominences, and coronal holes arise from (locally) open magnetic field lines.

High speed solar wind streams are formed by higher speed solar wind originating from **coronal holes**. Higher speed streams are less tightly wound in the Parker spiral compared to slower ones, and at various distances the faster solar wind **overtakes** the slower wind ahead of it.




FIG. 4.13. Geometry of the interaction between fast solar wind (on less tightly wound spiral streamlines) and ambient solar wind (on more tightly wound spiral streamlines). The plasma is compressed where streamlines converge. (From Pizzo, 1985.)

What are high speed streams? **High speed solar wind streams** are formed by higher speed solar wind originating from **coronal holes**. Higher speed streams are less tightly wound in the Parker spiral compared to slower ones, and at various distances the faster solar wind **overtakes** the slower wind ahead of it.

This figure (from Introduction to space physics) shows a schematic of the interaction between parcels of fast and slow solar wind in the Parker spiral. The interaction forms compression regions where the fast solar wind catches up to the slow solar wind streams, and leaves a rarefaction region behind it.

A **stream interaction region** (SIR) forms at the compressed boundary between the fast and slow solar wind in a high speed stream. High speed streams from persistent coronal holes over multiple solar rotations are called **corotating interaction regions** (CIRs).




FIG. 4.13. Geometry of the interaction between fast solar wind (on less tightly wound spiral streamlines) and ambient solar wind (on more tightly wound spiral streamlines). The plasma is compressed where streamlines converge. (From Pizzo, 1985.)

Now imagine a spacecraft sitting near the Earth in the previous diagram. What would it measure? These plots show an example of a **High speed solar wind stream** observed in-situ at ACE (at L1) using an ISWA layout.

The increase in speed from a solar wind high speed stream pumps energy into the magnetosphere which can cause **geomagnetic storms** and **energizes particles**.

They can produce energetic **electron flux enhancements** in the radiation belt.

Geomagnetic storms are disturbances/changes in Earth's magnetic field due to changes in solar wind conditions typically lasting 3-6 days.

High speed streams can also cause geomagnetic storms, however they are longer in duration and not as strong as geomagnetic storms caused by CMEs.

(the magnetosphere lesson will go into more detail)

How do high speed streams affect the Earth's magnetosphere?

The increase in speed from a solar wind high speed stream pumps energy into the magnetosphere which can cause **geomagnetic storms** and **energizes particles**.


They can produce energetic **electron flux enhancements** in the radiation belt.

Geomagnetic storms are disturbances/changes in Earth's magnetic field due to changes in solar wind conditions typically lasting 3-6 days.

High speed streams can also cause geomagnetic storms, however they are longer in duration and not as strong as geomagnetic storms caused by CMEs.

(the magnetosphere lesson will go into more detail)

A high speed stream can cause an energetic **electron flux enhancement** in the radiation belt.



These graphs show the same period in June 2012 when the high speed stream shown in the previous slides, caused an energetic **electron flux enhancement** in the radiation belt.

Slide link summary

SW REDI website

<http://ccmc.gsfc.nasa.gov/support/SWREDI/swredi.php>

iSWA <http://iswa.gsfc.nasa.gov>

iSWA Cygnet Glossary http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Full_iSWA_Cygnet_List

iSWA Space Weather Glossary <http://iswa3.ccmc.gsfc.nasa.gov/wiki/index.php/Glossary>

Most figures and tables in the slides are from the "Introduction to Space Physics" textbook

<http://www.cambridge.org/us/knowledge/isbn/item1145043>

iSWA layout of a high speed stream observed at ACE <http://go.nasa.gov/17nkicp>