







# Inner heliosphere modeling with LFM and Gamera

Space Plasma Theory and Modeling @ APL & collaborators

S. Merkin (JHU/APL)  
K. Sorathia (JHU/APL)  
L. Daldorff (JHU/APL)  
B. Zhang (HKU)  
J. Lyon (Dartmouth)  
M. Wiltberger (NCAR)


# APL space weather simulation toolbox






# LFM-helio code

- LFM — MHD code developed by J. Lyon, J. Fedder and C. Mobarry at NRL in the 80's
- Mainly applied to terrestrial magnetosphere; modified for inner heliosphere (Merkin et al., 2011, 2016a,b); regional plasma problems (Merkin et al., 2015); planetary magnetospheres (Zhang et al., 2018)
- LFM numerics
  - Handles arbitrary, non-orthogonal (and singular!) grids
  - High-order spatial reconstruction
  - Can capture 3D dynamics with relatively few cells
  - Intrinsically div-free B updates (constrained transport)
- Rotating/inertial frame calculations
- Time-dependent coronal boundary conditions



# LFM-helio: Inner heliosphere modeling




- LFM serves as backbone for the inner heliosphere modeling pipeline
- Modeling is time-dependent
- ADAPT/WSA-driven for quiet heliosphere
- Coupled with MAS for CME propagation



# Time-dependent quiet heliosphere

Driven by ADAPT/WSA

@ 0.1 AU



- ADAPT-driven WSA solutions
- 2-h to 1-d cadence
- Provide time-dependent inputs into MHD models of the solar wind
- Major problem: Radial magnetic field boundary condition should guarantee


$$\nabla \cdot \mathbf{B} = 0$$

Merkin et al., JGR, [2016]

# Time-dependent quiet heliosphere

Driven by ADAPT/WSA

@ 0.1 AU



$$\mathbf{E}_{\perp} = \nabla_{\perp} \times \hat{\mathbf{r}} \Psi$$



$$\Delta_{\perp} \Psi = \frac{\partial B_r}{\partial t}$$

From  
ADAPT/WSA



$$\mathbf{E}_{\perp}$$

MHD boundary  
condition

Merkin et al., JGR, [2016]  
Linker et al., AIP, [2016]





# Time-dependent quiet heliosphere

Driven by ADAPT/WSA

● ACE ● STEREO A ● STEREO B ● MESSENGER



- HCS moves around
- Transient SW velocity streams
- Complex HCS crossings/transitions



Merkin et al., JGR, 2016

# Time-dependent quiet heliosphere

Driven by ADAPT/WSA






Merkin et al., JGR, 2016




# Time-dependent heliosphere: Coronal mass ejections

MAS-driven CME propagation: First-principles from eruption to Earth



# Time-dependent heliosphere: Coronal mass ejections

MAS-driven CME propagation: First-principles from eruption to Earth



LFM-helio driven by the MHD-around-a-sphere (MAS) code.  
Work done in collaboration with Predictive Science Inc.

Merkin et al., ApJ, 2016




# Going forward

- LFM underlying code was written a while ago (30-40 years)
- Optimized for architectures long gone
- The code is robust and performs well on existing architectures
- Speed and portability becoming an issue with time
- LFM numerics are quite unique
  - Handles arbitrary, non-orthogonal (and singular!) grids
  - High-order spatial reconstruction
  - Can capture 3D dynamics with relatively few cells
  - Intrinsically div-free B updates (constrained transport)
- Needs a serious upgrade for next generation supercomputers


## What's next?

# Gamera


- Gamera is a reinvention of LFM
- Built to tackle modern challenges, but preserve the unique numerics of LFM
- Code written to expose multiple layers of heterogeneous parallelism
- Fortran 2003+, minimal external library dependence
- Portable, user-friendly, flexible
- Model coupling for inter-connected, multi-physics, multi-scale systems
- Standard tests to be published
- Website: [civspace.jhuapl.edu/gamera](http://civspace.jhuapl.edu/gamera)




DB: Data\_1250.vtk  
Cycle: 12500 Time: 0.997035




user: scrofula  
Wed Nov 9 10:49:16 2016



Hydro Kelvin-Helmholtz  
Elapsed Time: 5.05 (s)



Saturn magnetosphere



Inner heliosphere

# Conclusions

- Diverse set of space weather simulation tools
- Primary applications:
  - Geospace
    - Magnetosphere-ionosphere-thermosphere coupling
    - Radiation belts
    - Ring current
  - Inner heliosphere
- Backbone is the global magnetosphere MHD code (LFM, and now Gamera)
- Developed time-dependent boundary conditions for LFM-helio, for coupling with ADAPT/WSA and MAS coronal codes
- Currently working on implementing the same in Gamera
- Including Gibson-Low flux ropes @ inner boundary (~0.1 AU)
- Developed next generation LFM (Gamera)
- Gamera preserves (and improves upon) all the high heritage LFM numerics
- Already running heliosphere simulations at unprecedented resolution
- Resolving “emergent” structure (CIR substructure?)
- Are we at the scale where we can resolve MHD instabilities in the solar wind?
- Website: [civspace.jhuapl.edu/gamera](http://civspace.jhuapl.edu/gamera)
- Looking forward to working with CCMC on making it available

**GAMERA**  
Grid Agnostic MHD for Extended Research Applications







JOHNS HOPKINS  
APPLIED PHYSICS LABORATORY